



Small Grain Production in Southwest Iowa



Nathan Mueller
Ph.D. CCA

**Cropping
Systems
Extension
Educator for
Saline,
Jefferson, and
Gage counties**

Introduction




Nathan Mueller, Ph.D., CCA

Cropping Systems Extension
Educator

nathan.mueller@unl.edu

402-821-2151

croptechcafe.org



Overview

- Needs assessment by growers
- Wheat production in the region
- Economics and weather
- Soil health aspect
- Winter wheat management for new growers:
 - Variety Selection
 - Diseases
 - Planting dates and rates
 - Nutrients





Needs Assessment

Needs Assessment

1. Variety trials

1. Added location in 2017
2. More changes this fall

2. Wheat in rotation research

1. On-farm research studies through NRCS Soil Health Demonstration Farms

3. Soil fertility research and recommendations

1. Nitrogen and sulfur fertility trials

Winter Wheat Production Needs Assessment in Eastern Nebraska


Eastern Nebraska grower (n=37) and UNL and NRCS faculty/staff in responses (n=7) conducted in 2015 and 2019.

- Continue variety trials for eastern Nebraska
 - No-till after soybeans and seed corn
- Wheat in a corn/soybean rotation research
 - Research to document the impact wheat in rotation with corn and soybeans has on reducing soil erosion
 - Research to document the economic and rotation benefits and profitability to adding wheat to the corn/soybean rotation such as those below:
 - Rainfed corn-soybean-wheat/cover crop (3 crops in 3 years)
 - Rainfed corn-soybean-wheat/forage crop (4 crops in 3 years)
 - Irrigated corn-soybean-wheat/double-crop soybean (4 crops in 3 years)
 - Has this changed with climate change/growing season length?
 - BMPs for system and modeling to assess risk of not reaching maturity
 - Irrigated seed corn or short season corn-wheat-double crop soybeans (3 crops in 2 years)
- Soil fertility research and current fertility programs
 - Evaluate the need for chloride and sulfur applications in various soils
- Row spacing and seeding rate
 - Diseases risk and yield
- Predicting lodging risk and mitigation
 - Palisade (growth regulator), variety, N rate
- Cover crops after wheat research
 - Species, mix, management, etc. to benefit the next crops and economics
- On-farm storage BMPs and marketing considerations/plans for both grain and straw
- Managing winter wheat for cover crop seed production
- Crop rotation considerations
 - Cover crop after wheat for grazing or nitrogen credit
 - Mix versus mono-legume
 - Soil type consideration for cover crop
 - Corn, soybeans or other crops following year after wheat
 - After seed corn versus after soybeans
 - Shorter season corn and soybeans, potential yield penalty
- High yield practices
 - Plant population/stand counts
- Fungicide application methods and volume




Wheat Production in the Region

U.S. Wheat Production and Classes




● HARD RED WINTER ● HARD RED SPRING ● SOFT RED WINTER ● SOFT WHITE ● HARD WHITE ● DURUM


CropScape Wheat Frequency – MODIS



Retreat of Wheat in the East



Soil Moisture Regime - NRCS






Economics to Consider


USDA-NASS Winter Wheat Yield Trends

Winter Wheat Yields for Crop Reporting Districts




Local Success Stories

- Growing 100 bushel/acre wheat not uncommon
 - Thurston, Washington, and Jefferson counties
- UNL Variety Trials – Washington County
 - 2019 = 123 bpa (Plot average = 106)
 - 2018 = 110 bpa (Plot average = 99)
 - 2017 = 140 bpa (Plot average = 111)




Markets

- Grain - Strong Local Basis
 - Scoular Fremont = +0.30
 - ADM Lincoln = +0.30
- Straw – Rock Valley Hay Auction
 - Large Rounds - \$70 to \$135 per ton
 - Going rate in my areas was \$100 per ton




150 mile radius from Fremont

Corn Yields in w/Wheat in Rotation



Soybean Yields w/Wheat in Rotation

Monmouth 1998-2014






The Niche for Winter Wheat

Winter Wheat Can Help

- Manure management flexibility
- Consider all potential profits streams
 - Value of straw
 - Double crop and forages crop options
 - Potential nitrogen credit for legume cover crop
 - Corn and soybean yield improvement in 3-yr rotation
 - EQIP and CSP opportunities
- Weed control & herbicide cost
 - Marestalk, Palmer Amaranth, & Waterhemp
- Soil health and conservation
 - Soil structure, erosion control, cost-share and priority
- Workload management
- Manage weather risks



Managing Risks from Extreme Weather




Seasonal crop water use (ET) in Eastern Nebraska when water is not limiting.


Crop	Inches/year
Corn	21-24
Soybean	20-22
Winter Wheat	16-18
Alfalfa	31-35

Source: water.unl.edu

Precipitation Trends: Southeast NE




Precipitation Trends: Southeast NE





Wheat, Your Underground Partner

Water Stable Aggregates



Crop rotation and tillage matter



Improving Soil Health with Winter Wheat

- Aspects of soil health
 - Physical
 - Biological
 - Chemical

Improving Soil Health with Winter Wheat



Current issue

Soil health (biological, physical, and chemical) has been a popular focus over the past decade with emphasis on utilizing no-till and cover crops in eastern Nebraska. However, a more diverse crop rotation is often left out of the discussion as a way to improve soil health. The corn-soybean rotation is the most widely utilized cropping systems in eastern Nebraska. Despite the potential benefits adding a third or fourth crop to this rotation to improve soil health, few farmers in eastern Nebraska have because of various adoption barriers.

Crop rotation research

Two long-term (14 & 15 years) crop rotation studies in the Midwest have shown that including winter wheat into the corn-soybean rotation results in the following improvement in soil health:

1. Increase in water stable aggregates (most sensitive and best single indicator of soil physical health)
2. Higher total nitrogen
3. Higher potentially mineralizable nitrogen

These aspects of soil health were increased by adding wheat into the rotation regardless of the tillage system, conventional and no-till. The dense fibrous root system of wheat and nitrogen derived from wheat root deposits is likely the cause of these measurable differences. These long-term crop rotations studies did not include cover crops.

Overcome barriers to adding wheat

Barriers producers share

- Economics of wheat grain yield only/input cost compared to corn and soybean production
- Logistics of planting and harvesting only 1 or 2 fields
- Concerns about the learning curve of growing a new crop

Overcoming these barriers

- **Improve economics** by capturing good basis in Fremont, selling straw, growing forage crop after wheat, higher soybean yield in rotation, and USDA programs
- **Improve logistics** with custom drilling and harvesting and business opportunity
- **Reduce learning curve** through new website, grower group email list, and future peer-learning group, and work with cropping systems extension educators

Local Grower quotes

"It (wheat) breaks up our corn-bean rotation and it creates more organic matter, because of the root mass."
Local Farmer – Fremont Tribune

"Winter wheat gives you an additional 45 to 60 days for the cover crop to grow, which results in more material to graze if you choose to, and more root mass to help build organic matter in the soil." Local Farmer - Nebraska Farmer Magazine

"I'm hoping the wheat residue will double as a mat for the soil and a cover crop to help suppress weeds as well. It might decrease some chemical costs." Local Farmer - Nebraska Farmer Magazine

For more information

Nathan Mueller, PhD, CCA
Nebraska Extension Cropping Systems Educator
402-727-2775 or nathan.mueller@unl.edu

Wheat resources for eastern Nebraska at croptechcafe.org/winterwheat



Winter Wheat Management for New Growers

Management Factors of Winter Wheat in Eastern Nebraska

Rank	Management Factor	Yield Difference (bu/ac)
1	Variety Selection	21
2	Foliar Fungicide	16
3	Row Spacing, 15 vs 7.5"	16
4	Planting Date (2 wks)	10
5	Nutrient Mgnt (30 lbs N)	7
6	Seeding Rate	5



Variety Selection

Variety Selection is Critical


Variety	3-year yield average (bu/ac)
WB4303	99
AM Eastwood	78

UNL Variety Trial Test Results

<http://cropwatch.unl.edu/winter-wheat-variety-test-results>

Colorado Wheat Variety Database

<http://ramwheatdb.com/database.php>



Courtesy of Laura Thompson, UNL

Variety Testing Locations – Eastern Nebraska



- Soils and practices
 - Washington
 - Moody/Marshall series
 - No-till after soybeans
 - Saunders
 - Filbert - very deep, somewhat poorly drained, very slowly permeable soils formed in loess
 - Tilled after oats
 - Lancaster
 - Crete - very deep, moderately well drained soils formed in loess
 - Tilled after oats

Wheat Seed Brands in Eastern Nebraska

- Husker Genetics (UNL)
- NuPride (UNL)
- Kansas Wheat Alliance (KSU)
- WestBred (Bayer)
- Limagrain Cereal Seeds
- AgriMaxx
- AgriPro (Syngenta)
- Dyna-Gro

Digital Variety Tour - <http://croptechcafe.org/winterwheat/>

Example: Zenda – Kansas Wheat Alliance Released 2016



Placement

- All counties

Highlights

- Great yield performance record in UNL 3-yr trials
- Moderate resistance to Fusarium Head Blight
- Excellent test weight

Management Suggestions

- Replacement for Everest
- Average drought tolerance

View at croptechcafe.org/winterwheat



Disease Management

Bacterial Leaf Streak/Black Chaff



- Disease
 - Caused by bacteria *Xanthomonas translucens* pv. *undulosa*
 - Overwinters in residue/soil
- Management
 - Pathogen free seed
 - Variety selection that are less susceptible (i.e. SY Wolf, Overland)


Fusarium Head Blight (Scab)



- Disease
 - Caused by fungus *Fusarium graminearum* & *boothii*
 - Reported in NE since 1898
 - Survives on residue, in soil
 - Rainfall/humidity during flowering
 - Vomitoxin (DON)
- Management
 - Crop rotation (C-S-A)
 - Variety selection for resistance
 - Disease risk mapping
 - Scouting
 - Foliar fungicide at early flowering

Fungicides for FHB

- Prosaro 421 SC (Bayer)
 - Prothioconazole (3, Triazole)
 - Tebuconazole (3)
- Caramba (BASF)
 - Metconazole (3)
- Miravis Ace (Syngenta)
 - Two modes of action
 - Propiconazole (3)
 - Pydiflumetofen (7, SDHI)



Varieties with Moderate Resistance for FHB

- Varieties in the Trials
 - WB4699 (3)
 - Overland (3)
 - SY Benefit (4)
 - Zenda (5)
 - WB4269 (5)
 - LCS Valiant (2019)



Leaf Rust



- Disease
 - Caused by fungus *Puccinia triticina*
 - Does not overwinter
 - Central and eastern Nebraska
 - Yield losses up to 14% typical
- Management
 - Variety selection for resistance
 - Scouting
 - Foliar fungicide


Stripe Rust



- Disease
 - Caused by fungus *Puccinia striiformis* f. sp. *Tritica*
 - Does not overwinter
 - Has become a significant disease since 2010 in Nebraska
 - Yield loss up to 40%
- Management
 - Variety selection for resistance
 - Scouting
 - Foliar fungicide

Foliar Fungicide at Flag Leaf


Saunders County: Variety Millennium with Quilt



Wegulo, S. N., Zwingman, M. V., Breathnach, J. A. & Baenziger, P. S. 2011. Economic returns from fungicide application to control foliar fungal diseases in winter wheat. *Crop Protection*, Vol. 30, No. 6, (April 2011), pp. 685-692, ISSN 0261-2194

Foliar Fungicide at Flag Leaf

Lancaster County: Average across 6 varieties




Bhatta, M. 2015. Effect of genotype, environment, and production packages on yield, agronomic characteristics, and end-use quality of winter wheat. Master's thesis, Univ. of Nebraska, Lincoln, NE. <http://digitalcommons.unl.edu/agronhortdiss/98>.




Planting Date and Seeding Rate

Nebraska East District for 2006-2017



Planting Dates – Eastern Districts


- Recommended
 - Sept. 25 to Oct. 15
- Crop Insurance
 - 100% coverage until Oct. 10 or 15
- Late planting
 - Until Oct. 25 or 30



Target Planting Date Map


Target planting dates

- Planting date allows for 400 GDU (Base 40) accumulation between planting and December 31
- 1980s work in southeast NE
- Temperature norms – 1981-2010 used




Planting Dates - Iowa

Iowa



Wheat - Winter



Commodity - 11

Insurance Plan:
 1 - Yield Protection
 2 - Revenue Protection
 3 - Rev Protect w/ Harvest Exclusion

Saint Paul Region 2018 Final Plant Dates

Federal Crop Insurance

No Program

Planted acreage in a "No Program" county may be insurable by written agreement.



USDA Risk Management Agency

Map Creation Date: February 14, 2018

Map Projection: Web Mercator

RMA Business Analytics Division

The information displayed in this map is intended as an aid in displaying data provided or stored by the Risk Management Agency. It does not modify, replace or supersede any USDA published policy provisions or procedures. USDA Risk Management Agency's 2018 crop year sales, planting and reporting dates are published on the County Actuarial Table Special Provisions of Insurance document in the Actuarial Information Browser at <http://webapp.rma.usda.gov/apps/actuarialinformationbrowser2018/CropCriteria.asp>



Use Certified Seed

How to Read a Certified Analysis Tag

PVP STATEMENT
This variety may only be sold as certified seed. Any other transfer or sale of this seed is prohibited by federal law.

CERTIFIED SEED TAG
The blue certified tags assure the buyer is getting quality certified seed.

GERMINATION
The percentage of pure seed that will germinate in a controlled lab environment.

LOT NUMBER
Each lot of seed has a unique number that appears on all documents so that it can be traced back to the field where it was grown.

ORIGIN
Origin is the state where the seed was grown.

BRAND NAME
HUSKER GENETICS OVERLAND BRAND

VARIETY NAME
NEO1643 HRW WHEAT

PURE SEED
The percentage of weight of seed that is the named species.

CROP SEED
The percentage of seeds by weight that is other than the named species not considered weeds. Crop seed must be listed by name if over 5 percent.

INERT MATTER
The percentage by weight of material that will not grow.

WEED SEED
The percentage of weed seeds in this lot.

NOXIOUS WEEDS
The amount of seeds of weeds prohibited by state law.

TEST DATE
The month and year this lot was lab tested.

TOTAL VIABLE
Germination percentage plus the hard or dormant seed.

ANALYSIS TAG CONTENTS:

CERTIFIED SEED

UNAUTHORIZED PROPAGATION PROHIBITED. U.S. PLANT VARIETY PROTECTED. PVP 1994

HUSKER GENETICS OVERLAND BRAND

NEO1643 HRW WHEAT

PURE SEED %	99.50	GERMINATION %	90
CROP SEED %	.00	HARD/DORMANT %	0
INERT MATTER %	0.50	TOTAL VIABLE %	90
WEED SEED %	.00	TEST DATE	08/13
NOXIOUS WEED SEEDS NONE PER LB.			

LOT#W0-428

ORIGIN: NE

12156 SEEDS/LB


LBS NET WT

Quality certified by NEBRASKA CROP IMPROVEMENT ASSOCIATION
MEMBER OF ASSOCIATION OF OFFICIAL SEED CERTIFYING AGENCIES

* The Certifying Agency makes no warranty of any kind, express or implied, including merchantability or fitness for purpose, or otherwise, which extends beyond the certification that the seeds inspected met the regulations of this agency. The Seller guarantees this seed to conform to the analysis shown. No further warranty is expressed or implied. Sellers liability is limited to the purchase price of the seed.

Seeding Rate

Lancaster County, planted Oct. 2, 2013



Bhatta et al., 2017. Seeding rate, genotype, and top-dressed nitrogen effect on yield and agronomic characteristics of winter wheat. *Crop Sci.* 57:951-963

Planting Date by Seeding Rate




Figure 2. Irrigated wheat yields from 2017 for a study conducted in northwest Kansas evaluating planting dates and seeding rates.

Seeding Rate and Planting Date

Crop Tech Cafe Winter Wheat Seeding Rate Calculator for East Central Nebraska		
Estimated Planting Dates	Oct. 7 - Oct. 14	Select estimated planting dates from drop-down
Recommended Seeding Rate (Pure Live Seeds per Acre)	1,350,000	Based on estimated planting dates selected
Germination (%)	95	Enter germination from seed tag
Purity (%)	99	Enter purity from seed tag
Seed Size (seeds/lbs)	15,000	Enter seed size from seed tag
Contact Nathan Mueller at nathan.mueller@unl.edu with questions	Seeding Rate (lbs/acre)	96
	Seeding Rate (bu/acre)	1.6

- Plant certified fungicide-treated seed to control seed-transmitted and soilborne fungal diseases
- Plant at 1.5 inches deep no-till after soybeans

Download at croptechcafe.org/winterwheat



Evaluating Winter Wheat Stands

- Better option than a tape measure
- No need to glue together, so easy storage.
- Interpretations
 - Assesses yield components
 - Risk of weed pressure

Evaluating Winter Wheat Stands



Making your own grid for 1/10,000 of an acre

Supplies:

- One 10 ft piece of ½ inch PVC & Four PVC elbows for ½ inch pipe
- Measuring tape, saw, & permanent marker

Cut and Assemble: Dimensions to cut pipe for each row spacing:

- 7.5 inch row spacing
 - 22.5" wide and 28" long
- 8 inch row spacing
 - 24" wide and 26" long
- 10 inch row spacing
 - 30" wide and 21" long

Mark: With your permanent marker, add 3 lines (7.5, 8, or 10) inches apart to represent rows on each two sides (the side that is 22.5, 24, or 30" wide).

Assess your winter wheat stand

Count the number of plants for the three rows inside the grid in the fall or early spring with a tally counter (shown in picture) in several different areas of the field and then average those values. Interpretation of your counts from 1/10,000 of an acre grid:

- **Less than 50 plants** – Likely due to high variability in the stand, consider replanting at an angle or using as a cover crop
- **50 to 65 plants** – Reduced yield, can still yield well with good tillering and weed control
- **65 to 95 plants** – Good, can obtain close to max yield potential
- **More than 95 plants** – Ideal

For more information

Nathan Mueller, PhD, CCA
Nebraska Extension Cropping Systems Educator
402-727-2775 or nathan.mueller@unl.edu

Current issue

Recommended seeding rates start at 1.2 million seeds per acre and increase as planting is delayed through October in eastern Nebraska. You can download the seeding rate Excel tool at croptechcafe.org/winterwheat Evaluating your winter wheat stand and determining how many plants you have on a per acre basis is something most growers find difficult and time consuming. It is normal to use a tape measure to help count plants for corn and soybean, but not as ideal for wheat. Knowing what percentage of your seeding rate became established as plants is critical information to have to make future planting adjustments and also to make a determination, when stands are poor, to use the wheat as a cover crop.

Wheat resources for eastern Nebraska at croptechcafe.org/winterwheat

N P K S Ca Mg

Nutrient Management

Fe Mn Cu Zn B Cl Mo Ni

Nitrogen Management

- Based on regional UNL recommendations and local grower experiences
 - 80 – 110 lbs N/acre
- Apply most or all as wheat begins to green up in February/March
- Grain protein can be improved with late N applications
 - N at Flag leaf can still improved protein/yield

A large, bold, red letter 'N' with a slight shadow effect, positioned on the right side of the slide.

Phosphorus Management

- Higher soil test phosphorus needed compared to corn and soybeans
 - Similar to alfalfa and corn-after-corn
 - 25 ppm Bray P1 or more
- Helps with early growth, tillering, and winter hardiness

P

Sulfur Management

- Becoming more common in northeast and north central KS and southeast NE
- No-till, cold springs, low organic matter eroded areas
- Pale yellow color in the spring, entire plant, more so on younger leaves
- 10 to 20 lbs S/acre
- Dry ammonium sulfate or liquid ammonia thiosulfate



Photo by Stephen Wegulo, UNL

Chloride Management

- Recent analysis in Kansas performed across multiple years and locations suggest an average yield response of 8% to chloride fertilization.
- Chloride has been shown to suppress take-all root rot, tan spot, stripe rust, leaf rust, and Septoria
- Research has shown equal performance to both preplant and topdress applications.



Table 5. *Soil test chloride interpretation and fertilizer recommendation.*

Category	Soil Chloride*		Cl Recommended
	lb/a	ppm	lb/a
Low	< 30	< 4	20
Medium	30-45	4-6	10
High	> 45	> 6	0

**Interpretations valid for 0-24 inch samples on wheat, corn and grain sorghum.*



Resources

Wheat Wheat Cafe




Welcome to the Winter Wheat Cafe for Eastern Nebraska

Positioning your farm to manage manure, control tough weeds, and improve soil health are just some of the advantages to growing winter wheat in Eastern Nebraska. Current livestock producers and future poultry growers will find additional value of adding winter wheat to their farming operation.


Resources found on this page:

application timing (select all that apply)? *Split application (preplant and sidedress)* (44%, 11

Votes)



Spring liquid (UAN or liquid mixes) (24%, 6 Votes)



Split application (preplant and fertigation) (16%, 4 Votes)



Fall anhydrous ammonia (12%, 3 Votes)



Spring anhydrous ammonia (4%, 1 Votes)



Spring dry (Urea, ammonium nitrate, etc.) (0%, 0 Votes)



Total Voters: 25

Brought to You By




Serving Northeast Nebraska



View at croptechcafe.org/winterwheat



Hard Copy Resources



Feekes scale for cereal growth stages

SEEDLING GROWTH	1 One shoot, first leaf through coleoptile
TILLERING	2 Tillering begins; main shoot and one tiller
	3 Tillers formed; leaves often twisted In some varieties, plant may be prostrate in appearance
	4 Leaf sheaths lengthen; beginning pseudostem erection
	5 Leaf sheaths fully elongated to form strongly erect pseudostem
	6 First node of stem visible at base of shoot; jointing
STEM EXTENSION	7 Second node of stem formed; next-to-last leaf just visible
	8 Flag leaf visible but still rolled up
	9 Ligule of flag leaf just visible
	10 Flag leaf sheath completely grown out; booting
HEADING	10.1 First awns of head just visible
	10.2 1/4 of heading process complete
	10.3 1/2 of heading process complete
	10.4 3/4 of heading process complete
	10.5 All heads out of sheath
FLOWERING	10.5.1 Beginning of flowering
	10.5.2 Flowering complete to top of head
	10.5.3 Flowering complete at base of head
	10.5.4 Flowering complete; kernel watery ripe
RIPENING	11.1 Kernel milky ripe; milk stage
	11.2 Kernel mealy ripe; soft but dry consistency; soft dough stage
	11.3 Kernel hard; difficult to divide with thumbnail; hard dough stage
	11.4 Kernel harvest ready; straw dead



croptechcafe.org/winterwheat



[@croptechcafe](https://twitter.com/croptechcafe)



nathan.mueller@unl.edu

Thank You!